

LANDEN SURVEY

DANTE V. MANNA AND VICTOR H. MOLL

ABSTRACT. Landen transformations are maps on the coefficients of an integral that preserve its value. We present a brief survey of their appearance in the literature.

To Henry, who provides inspiration, taste and friendship

1. IN THE BEGINNING THERE WAS GAUSS

In the year 1985, one of us had the luxury of attending a graduate course on *Elliptic Functions* given by Henry McKean at the Courant Institute. Among the many beautiful results he described in his unique style, there was a calculation of Gauss: take two positive real numbers a and b , with $a > b$, and form a new pair by replacing a with the arithmetic mean $(a + b)/2$ and b with the geometric mean $\sqrt{ab}$. Then iterate:

$$(1.1) \quad a_{n+1} = \frac{a_n + b_n}{2}, \quad b_{n+1} = \sqrt{a_n b_n}$$

starting with $a_0 = a$ and $b_0 = b$. Gauss [29] was interested in the initial conditions $a = 1$ and $b = \sqrt{2}$. The iteration generates a sequence of algebraic numbers which rapidly become impossible to describe explicitly, for instance,

$$(1.2) \quad a_3 = \frac{1}{2^3} \left((1 + \sqrt[4]{2})^2 + 2\sqrt{2}\sqrt[8]{2}\sqrt{1 + \sqrt{2}} \right)$$

is a root of the polynomial

$$\begin{aligned} G(a) = & 16777216a^8 - 16777216a^7 + 5242880a^6 - 10747904a^5 \\ & + 942080a^4 - 1896448a^3 + 4436a^2 - 59840a + 1. \end{aligned}$$

The numerical behavior is surprising; a_6 and b_6 agree to 87 digits. It is simple to check that

$$(1.3) \quad \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n.$$

See (6.1) for details. This common limit is called the *arithmetic-geometric mean* and is denoted by $\text{AGM}(a, b)$. It is the explicit dependence on the initial condition that is hard to discover.

Gauss computed some numerical values and observed that

$$(1.4) \quad a_{11} \sim b_{11} \sim 1.198140235$$

Date: February 1, 2008.

Key words and phrases. Integrals, arithmetic-geometric mean, elliptic integrals.

and then he *recognized* the reciprocal of this number as a numerical approximation to the elliptic integral

$$(1.5) \quad I = \frac{2}{\pi} \int_0^1 \frac{dt}{\sqrt{1-t^4}}.$$

It is unclear to the authors how Gauss recognized this number - he simply knew it. (Stirling's tables may have been a help; [12] contains a reproduction of the original notes and comments.) He was particularly interested in the evaluation of this definite integral as it provides the length of a lemniscate. In his diary Gauss remarked, '*This will surely open up a whole new field of analysis*' [27, 15].

Gauss' procedure to find an analytic expression for $\text{AGM}(a, b)$ began with the elementary observation

$$(1.6) \quad \text{AGM}(a, b) = \text{AGM}\left(\frac{a+b}{2}, \sqrt{ab}\right)$$

and the homogeneity condition

$$(1.7) \quad \text{AGM}(\lambda a, \lambda b) = \lambda \text{AGM}(a, b).$$

He used (1.6) with $a = (1 + \sqrt{k})^2$ and $b = (1 - \sqrt{k})^2$, with $0 < k < 1$, to produce

$$\text{AGM}(1 + k + 2\sqrt{k}, 1 + k - 2\sqrt{k}) = \text{AGM}(1 + k, 1 - k).$$

He then used the homogeneity of AGM to write

$$\begin{aligned} \text{AGM}(1 + k + 2\sqrt{k}, 1 + k - 2\sqrt{k}) &= \text{AGM}((1 + k)(1 + k^*), (1 + k)(1 - k^*)) \\ &= (1 + k) \text{AGM}(1 + k^*, 1 - k^*), \end{aligned}$$

with

$$(1.8) \quad k^* = \frac{2\sqrt{k}}{1 + k}.$$

This resulted in the functional equation

$$(1.9) \quad \text{AGM}(1 + k, 1 - k) = (1 + k) \text{AGM}(1 + k^*, 1 - k^*).$$

In his analysis of (1.9), Gauss substituted the power series

$$(1.10) \quad \frac{1}{\text{AGM}(1 + k, 1 - k)} = \sum_{n=0}^{\infty} a_n k^{2n}$$

into (1.9) and, solved an infinite system of nonlinear equations, to produce

$$(1.11) \quad a_n = 2^{-2n} \binom{2n}{n}^2.$$

Then he recognized the series as that of an elliptic integral to obtain

$$(1.12) \quad \frac{1}{\text{AGM}(1 + k, 1 - k)} = \frac{2}{\pi} \int_0^{\pi/2} \frac{dx}{\sqrt{1 - k^2 \sin^2 x}}.$$

This is a remarkable tour de force.

The function

$$(1.13) \quad K(k) = \int_0^{\pi/2} \frac{dx}{\sqrt{1 - k^2 \sin^2 x}}$$

is the *elliptic integral of the first kind*. It can also be written in the algebraic form

$$(1.14) \quad K(k) = \int_0^1 \frac{dt}{\sqrt{(1-t^2)(1-k^2t^2)}}.$$

In this notation, (1.9) becomes

$$(1.15) \quad K(k^*) = (1+k)K(k).$$

This is the *Landen transformation* for the complete elliptic integral. John Landen [37], the namesake of the transformation, studied related integrals: for example,

$$(1.16) \quad \kappa := \int_0^1 \frac{dx}{\sqrt{x^2(1-x^2)}}.$$

He derived identities such as

$$(1.17) \quad \kappa = \varepsilon \sqrt{\varepsilon^2 - \pi}, \quad \text{where } \varepsilon := \int_0^{\pi/2} \sqrt{2 - \sin^2 \theta} \, d\theta,$$

proven mainly by suitable changes of variables in the integral for ε . In [52] the reader will find a historical account of Landen's work, including the above identities.

The reader will find in [15] and [43] proofs in a variety of styles. In trigonometric form, the Landen transformation states that

$$(1.18) \quad G(a, b) = \int_0^{\pi/2} \frac{d\theta}{\sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta}}$$

is invariant under the change of parameters $(a, b) \mapsto \left(\frac{a+b}{2}, \sqrt{ab}\right)$. D. J. Newman [46] presents a very clever proof: the change of variables $x = b \tan \theta$ yields

$$(1.19) \quad G(a, b) = \frac{1}{2} \int_{-\infty}^{\infty} \frac{dx}{\sqrt{(a^2 + x^2)(b^2 + x^2)}}.$$

Now let $x \mapsto x + \sqrt{x^2 + ab}$ to complete the proof. Many of the above identities can now be searched for and proven on a computer [12].

2. AN INTERLUDE: THE QUARTIC INTEGRAL

The evaluation of definite integrals of rational functions is one of the standard topics in Integral Calculus. Motivated by the lack of success of symbolic languages, we began a systematic study of these integrals. *A posteriori*, one learns that even rational functions are easier to deal with. Thus we start with one having a *power of a quartic* in its denominator. The evaluation of the identity

$$(2.1) \quad \int_0^\infty \frac{dx}{(x^4 + 2ax^2 + 1)^{m+1}} = \frac{\pi}{2^{m+3/2} (a+1)^{m+1/2}} P_m(a),$$

where

$$(2.2) \quad P_m(a) = \sum_{l=0}^m d_l(m) a^l$$

with

$$(2.3) \quad d_l(m) = 2^{-2m} \sum_{k=l}^m 2^k \binom{2m-2k}{m-k} \binom{m+k}{m} \binom{k}{l},$$

was first established in [7].

A standard hypergeometric argument yields

$$(2.4) \quad P_m(a) = P_m^{(\alpha, \beta)}(a),$$

where

$$(2.5) \quad P_m^{(\alpha, \beta)}(a) = \sum_{k=0}^m (-1)^{m-k} \binom{m+\beta}{m-k} \binom{m+k+\alpha+\beta}{k} 2^{-k} (a+1)^k$$

is the classical Jacobi polynomial; the parameters α and β are given by $\alpha = m + \frac{1}{2}$ and $\beta = -m - \frac{1}{2}$. A general description of these functions and their properties are given in [1]. The twist here is that they depend on m , which means most of the properties of P_m had to be proven from scratch. For instance, P_m satisfies the recurrence

$$\begin{aligned} P_m(a) &= \frac{(2m-3)(4m-3)a}{4m(m-1)(a-1)} P_{m-2}(a) - \frac{(4m-3)a(a+1)}{2m(m-1)(a-1)} P'_{m-2}(a) \\ &+ \frac{4m(a^2-1)+1-2a^2}{2m(a-1)} P_m(a). \end{aligned}$$

This *cannot* be obtained by replacing $\alpha = m + \frac{1}{2}$ and $\beta = -m - \frac{1}{2}$ in the standard recurrence for the Jacobi polynomials. The reader will find in [3] several different proofs of (2.1).

The polynomials $P_m(a)$ makes a surprising appearance in the expansion

$$(2.6) \quad \sqrt{a + \sqrt{1+c}} = \sqrt{a+1} \left[1 - \sum_{k=1}^{\infty} \frac{(-1)^k}{k} \frac{P_{k-1}(a) c^k}{2^{k+1} (a+1)^k} \right]$$

as described in [9]. The special case $a = 1$ appears in [22], page 191, exercise 21. Ramanujan [4] had a more general expression, but only for the case $c = a^2$:

$$(2.7) \quad (a + \sqrt{1+a^2})^n = 1 + na + \sum_{k=2}^{\infty} \frac{b_k(n) a^k}{k!},$$

where, for $k \geq 2$,

$$(2.8) \quad b_k(n) = \begin{cases} n^2(n^2-2^2)(n^2-4^2) \cdots (n^2-(k-2)^2) & \text{if } k \text{ is even,} \\ n(n^2-1^2)(n^2-3^2) \cdots (n^2-(k-2)^2) & \text{if } k \text{ is odd.} \end{cases}$$

This result appears in Berndt [4] as Corollary 2 to Entry 14 and is machine-checkable, as are many of the identities in this section.

The coefficients $d_l(m)$ in (2.3) have many interesting properties:

- They form a *unimodal sequence*: there exists an index $0 \leq m^* \leq m$ such that $d_j(m)$ increases up to $j = m^*$ and decreases from then on. See [6] for a proof of the more general statement: *If $P(x)$ is a polynomial with nondecreasing, nonnegative coefficients, then the coefficient sequence of $P(x+1)$ is unimodal.*
- They form a *log-concave sequence*: define the operator $\mathfrak{L}(\{a_k\}) := \{a_k^2 - a_{k-1}a_{k+1}\}$ acting on sequences of positive real numbers. A sequence $\{a_k\}$ is called log-concave if its image under $\mathfrak{L}$ is again a sequence of positive numbers; i.e. $a_k^2 - a_{k-1}a_{k+1} \geq 0$. Note that this condition is satisfied if and only if the sequence $\{b_k := \log(a_k)\}$ is

concave, hence the name. We refer the reader to [53] for a detailed introduction. The log-concavity of $d_l(m)$ was established in [35] using Computer Algebra techniques: in particular, cylindrical algebraic decompositions as developed in [24] and [26].

- They produce interesting polynomials: in [11] one finds the representation

$$(2.9) \quad d_l(m) = \frac{A_{l,m}}{l! m! 2^{m+l}},$$

with

$$(2.10) \quad A_{l,m} = \alpha_l(m) \prod_{k=1}^m (4k-1) - \beta_l(m) \prod_{k=1}^m (4k+1).$$

Here α_l and β_l are polynomials in m of degrees l and $l-1$, respectively. For example, $\alpha_1(m) = 2m+1$ and $\beta_1(m) = 1$, so that the coefficient of the linear term of $P_m(a)$ is

$$(2.11) \quad d_1(m) = \frac{1}{m! 2^{m+1}} \left((2m+1) \prod_{k=1}^m (4k-1) - \prod_{k=1}^m (4k+1) \right).$$

J. Little established in [40] the remarkable fact that the polynomials $\alpha_l(m)$ and $\beta_l(m)$ have all their roots on the vertical line $\operatorname{Re} m = -\frac{1}{2}$.

When we showed this to Henry, he simply remarked: *the only thing you have to do now is to let $l \rightarrow \infty$ and get the Riemann hypothesis*. The proof in [40] consists in a study of the recurrence

$$(2.12) \quad y_{l+1}(s) = 2s y_l(s) - (s^2 - (2l-1)^2) y_{l-1}(s),$$

satisfied by $\alpha_l((s-1)/2)$ and $\beta_l((s-1)/2)$. There is no Number Theory in the proof, so it is not likely to connect to the Riemann zeta function $\zeta(s)$, but one never knows.

The arithmetical properties of $A_{l,m}$ are beginning to be elucidated. We have shown that their 2-adic valuation satisfies

$$(2.13) \quad \nu_2(A_{l,m}) = \nu_2((m+1-l)_{2l}) + l,$$

where $(a)_k = a(a+1)(a+2)\cdots(a+k-1)$ is the Pochhammer symbol. This expression allows for a combinatorial interpretation of the block structure of these valuations. See [2] for details.

3. THE INCIPIENT RATIONAL LANDEN TRANSFORMATION

The clean analytic expression in (2.1) is not expected to extend to rational functions of higher order. In our analysis we distinguish according to the domain of integration: the finite interval case, mapped by a bilinear transformation to $[0, \infty)$, and the whole line. In this section we consider the definite integral,

$$(3.1) \quad U_6(a, b; c, d, e) = \int_0^\infty \frac{cx^4 + dx^2 + e}{x^6 + ax^4 + bx^2 + 1} dx,$$

as the simplest case on $[0, \infty)$. The case of the real line is considered below. The integrand is chosen to be even by necessity: *none of the techniques in this section work for the odd case*. We normalize two of the coefficients in the denominator in order to reduce the number of parameters. The standard approach for the

evaluation of (3.1) is to introduce the change of variables $x = \tan \theta$. This leads to an intractable trigonometric integral.

A different result is obtained if one first symmetrizes the denominator: we say that a polynomial of degree d is *reciprocal* if $Q_d(1/x) = x^{-d}Q_d(x)$, that is, the sequence of its coefficients is a palindrome. Observe that if Q_d is any polynomial of degree d , then

$$(3.2) \quad T_{2d}(x) = x^d Q_d(x) Q_d(1/x)$$

is a reciprocal polynomial of degree $2d$. For example, if

$$(3.3) \quad Q_6(x) = x^6 + ax^4 + bx^2 + 1.$$

then

$$\begin{aligned} T_{12}(x) &= x^{12} + (a+b)x^{10} + (a+b+ab)x^8 + \\ &+ (2+a^2+b^2)x^6 + (a+b+ab)x^4 + (a+b)x^2 + 1. \end{aligned}$$

The numerator and denominator in the integrand of (3.1) are now scaled by $x^6 Q_6(1/x)$ to produce a new integrand with reciprocal denominator:

$$(3.4) \quad U_6 = \int_0^\infty \frac{S_{10}(x)}{T_{12}(x)} dx,$$

where we write

$$S_{10}(x) = \sum_{j=0}^5 s_j x^{2j} \text{ and } T_{12}(x) = \sum_{j=0}^6 t_j x^{2j}.$$

The change of variables $x = \tan \theta$ now yields

$$(3.5) \quad U_6 = \int_0^{\pi/2} \frac{S_{10}(\tan \theta) \cos^{10}(\theta)}{T_{12}(\tan \theta) \cos^{12}(\theta)} d\theta.$$

Now let $w = \cos 2\theta$ and use $\sin^2 \theta = \frac{1}{2}(1-w)$ and $\cos^2 \theta = \frac{1}{2}(1+w)$ to check that the numerator and denominator of the new integrand,

$$(3.6) \quad S_{10}(\tan \theta) \cos^{10} \theta = \sum_{j=0}^5 s_j \sin^{2j} \theta \cos^{10-2j} \theta$$

and

$$\begin{aligned} T_{12}(\tan \theta) \cos^{12} \theta &= \sum_{j=0}^6 t_j \sin^{2j} \theta \cos^{12-2j} \theta \\ &= 2^{-6} \sum_{j=0}^6 t_j (1-w)^j (1+w)^{6-j}, \end{aligned}$$

are both polynomials in w . The mirror symmetry of T_{12} , reflected in $t_j = t_{6-j}$, shows that the new denominator is an *even* polynomial in w . The symmetry of cosine about $\pi/2$ shows that the terms with odd power of w have a vanishing integral. Thus, with $\psi = 2\theta$, and using the symmetry of the integrand to reduce the integral from $[0, \pi]$ to $[0, \pi/2]$, we obtain

$$U_6 = \int_0^{\pi/2} \frac{r_4 \cos^4 \psi + r_2 \cos^2 \psi + r_0}{q_6 \cos^6 \psi + q_4 \cos^4 \psi + q_2 \cos^2 \psi + q_0} d\psi.$$

The parameters r_j, q_j have explicit formulas in terms of the original parameters of U_6 . This even rational function of $\cos \psi$ can now be expressed in terms of $\cos 2\psi$ to produce (letting $\theta \leftarrow 2\psi$)

$$U_6 = \int_0^\pi \frac{\alpha_2 \cos^2 \theta + \alpha_1 \cos \theta + \alpha_0}{\beta_3 \cos^3 \theta + \beta_2 \cos^2 \theta + \beta_1 \cos \theta + \beta_0} d\theta.$$

The final change of variables $y = \tan \frac{\theta}{2}$ yields a new rational form of the integrand:

$$(3.7) \quad U_6 = \int_0^\infty \frac{c_1 y^4 + d_1 y^2 + e_1}{y^6 + a_1 y^4 + b_1 y^2 + 1} dy.$$

Keeping track of the parameters, we have established:

Theorem 3.1. *The integral*

$$(3.8) \quad U_6 = \int_0^\infty \frac{cx^4 + dx^2 + e}{x^6 + ax^4 + bx^2 + 1} dx$$

is invariant under the change of parameters

$$(3.9) \quad \begin{aligned} a_1 &\leftarrow \frac{ab + 5a + 5b + 9}{(a + b + 2)^{4/3}}, \\ b_1 &\leftarrow \frac{a + b + 6}{(a + b + 2)^{2/3}}, \end{aligned}$$

for the denominator parameters and

$$\begin{aligned} c_1 &\leftarrow \frac{c + d + e}{(a + b + 2)^{2/3}}, \\ d_1 &\leftarrow \frac{(b + 3)c + 2d + (a + 3)e}{a + b + 2}, \\ e_1 &\leftarrow \frac{c + e}{(a + b + 2)^{1/3}} \end{aligned}$$

for those of the numerator.

Theorem 3.1 is the precise analogue of the elliptic Landen transformation (1.1) for the case of a rational integrand. We call (3.9) a *rational Landen transformation*. This construction was first presented in [8].

3.1. Even rational Landen transformations. More generally, there is a similar transformation of coefficients for *any even rational integrand*; details appear in [10]. We call these *even rational Landen Transformations*. The obstruction in the general case comes from (3.6); one does not get a polynomial in $w = \cos 2\theta$.

The method of proof for even rational integrals can be summarized as follows.

1) Start with an even rational integral:

$$U_{2p} = \int_0^\infty \frac{\text{even polynomial in } x}{\text{even polynomial in } x} dx.$$

2) Symmetrize the denominator to produce

$$U_{2p} = \int_0^\infty \frac{\text{even polynomial in } x}{\text{even reciprocal polynomial in } x} dx.$$

The degree of the denominator is doubled.

3) Let $x = \tan \theta$. Then

$$U_{2p} = \int_0^{\pi/2} \frac{\text{polynomial in } \cos 2\theta}{\text{even polynomial in } \cos 2\theta} d\theta.$$

4) Symmetry produced the vanishing of the integrands with an odd power of $\cos \theta$ in the numerator. We obtain

$$U_{2p} = \int_0^{\pi/2} \frac{\text{even polynomial in } \cos 2\theta}{\text{even polynomial in } \cos 2\theta} d\theta.$$

5) Let $\psi = 2\theta$ to produce

$$U_{2p} = \int_0^\pi \frac{\text{even polynomial in } \cos \psi}{\text{even polynomial in } \cos \psi} d\psi.$$

Using symmetry this becomes an integral over $[0, \pi/2]$.

6) Let $y = \tan \psi$ and use $\cos \psi = 1/\sqrt{1+y^2}$ to obtain

$$U_{2p} = \int_0^\infty \frac{\text{even polynomial in } y}{\text{even polynomial in } y} dy.$$

The degree of the denominator is half of what it was in Step 5.

Keeping track of the degrees one checks that the degree of the new rational function is the same as the original one, with new coefficients that appear as functions of the old ones.

4. A GEOMETRIC INTERPRETATION

We now present a geometric foundation of the general even rational Landen transformation (3.9) using the theory of Riemann surfaces. The text [51] provides an introduction to this theory, including definitions of objects we will refer to here. The sequence of transformations in section 3 can be achieved in one step by relating $\tan 2\theta$ to $\tan \theta$. For historical reasons (this is what we did first) we present the details with *cotangent* instead of tangent.

Consider the even rational integral

$$(4.1) \quad I = \int_0^\infty R(x) dx = \frac{1}{2} \int_{-\infty}^\infty R(x) dx.$$

Introduce the new variable

$$(4.2) \quad y = R_2(x) = \frac{x^2 - 1}{2x},$$

motivated by the identity $\cot 2\theta = R_2(\cot \theta)$. The function $R_2 : \mathbb{R} \rightarrow \mathbb{R}$ is a two-to-one map. The sections of the inverse are

$$(4.3) \quad x = \sigma_\pm(y) = y \pm \sqrt{y^2 + 1}.$$

Splitting the original integral as

$$(4.4) \quad I = \int_{-\infty}^0 R(x) dx + \int_0^{\infty} R(x) dx$$

and introducing $x = \sigma_+(y)$ in the first and $x = \sigma_-(y)$ in the second integral, yields

$$(4.5) \quad I = \int_{-\infty}^{\infty} (R_+(y) + R_-(y)) dy$$

where

$$R_+(y) = R(\sigma_+(y)) + R(\sigma_-(y)) \text{ and } R_-(y) = \frac{y}{\sqrt{y^2 + 1}} (R(\sigma_+(y)) - R(\sigma_-(y))).$$

A direct calculation shows that R_+ and R_- are rational functions of degree at most that of R .

The change of variables $y = R_2(x)$ converts the meromorphic differential $\varphi = R(x) dx$ into

$$\begin{aligned} R(\sigma_+(y)) \frac{d\sigma_+}{dy} + R(\sigma_-(y)) \frac{d\sigma_-}{dy} &= \left((R(\sigma_+) + R(\sigma_-)) + \frac{y(R(\sigma_+) - R(\sigma_-))}{\sqrt{y^2 + 1}} \right) dy \\ &= (R_+(y) + R_-(y)) dy. \end{aligned}$$

The general situation is this: start with a finite ramified cover $\pi : X \rightarrow Y$ of Riemann surfaces and a meromorphic differential φ on X . Let $U \subset Y$ be a simply connected domain that contains no critical values of π , and let $\sigma_1, \dots, \sigma_k : U \rightarrow X$ be the distinct sections of π . Define

$$(4.6) \quad \pi_* \varphi \Big|_U = \sum_{j=1}^k \sigma_j^* \varphi.$$

In [33] we show that this construction preserves analytic 1-forms, that is, if φ is an analytic 1-form in X then $\pi_* \varphi$ is an analytic 1-form in Y . Furthermore, for any rectifiable curve γ on Y , we have

$$(4.7) \quad \int_{\gamma} \pi_* \varphi = \int_{\pi^{-1}\gamma} \varphi.$$

In the case of projective space, this leads to the following:

Lemma 4.1. *If $\pi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is analytic, and $\varphi = R(z) dz$ with R a rational function, then $\pi_* \varphi$ can be written as $R_1(z) dz$ with R_1 a rational function of degree at most the degree of R .*

This is the generalization of the fact that the integrals in (4.1) and (4.5) are the same.

5. A FURTHER GENERALIZATION

The procedure described in Section 3 can be extended with the rational map R_m , defined by the identity

$$(5.1) \quad \cot m\theta = R_m(\cot \theta).$$

Here $m \in \mathbb{N}$ is arbitrary greater or equal than 2. We present some elementary properties of the rational function R_m .

Proposition 5.1. The rational function R_m satisfies:

1) For $m \in \mathbb{N}$ define

$$P_m(x) := \sum_{j=0}^{\lfloor m/2 \rfloor} (-1)^j \binom{m}{2j} x^{m-2j} \text{ and } Q_m(x) := \sum_{j=0}^{\lfloor (m-1)/2 \rfloor} (-1)^j \binom{m}{2j+1} x^{m-(2j+1)}.$$

Then $R_m := P_m/Q_m$.

2) The function R_m is conjugate to $f_m(x) := x^m$ via $M(x) := \frac{x+i}{x-i}$; that is, $R_m = M^{-1} \circ f_m \circ M$.

3) The polynomials P_m and Q_m have simple real zeros given by

$$p_k := \cot\left(\frac{(2k+1)\pi}{2m}\right) \text{ for } 0 \leq k \leq m-1, \text{ and } q_k := \cot\left(\frac{k\pi}{m}\right) \text{ for } 1 \leq k \leq m-1.$$

If we change the domain to the entire real line, we can, using the rational substitutions $R_m(x) \mapsto x$, produce a rational Landen transformation for an arbitrary integrable rational function $R(x) = B(x)/A(x)$ for each integer value of m . The result is a new list of coefficients, from which one produces a second rational function $R^{(1)}(x) = J(x)/H(x)$ with

$$(5.2) \quad \int_{-\infty}^{\infty} \frac{B(x)}{A(x)} dx = \int_{-\infty}^{\infty} \frac{J(x)}{H(x)} dx.$$

Iteration of this procedure yields a sequence $\mathbf{x}_n$, that has a limit $\mathbf{x}_\infty$ with convergence of order m , that is,

$$(5.3) \quad \|\mathbf{x}_{n+1} - \mathbf{x}_\infty\| \leq C \|\mathbf{x}_n - \mathbf{x}_\infty\|^m.$$

We describe this procedure here in the form of an algorithm; proofs appear in [41].

Lemma 4.1 applied to the map $\pi(x) = R_m(x)$, viewed as ramified cover of $\mathbb{P}^1$, guarantees the existence of a such new rational function $R^{(1)}$. The question of effective computation of the coefficients of J and H is discussed below. In particular, we show that all these calculations can be done symbolically.

• Algorithm for Deriving Rational Landen Transformations

Step 1. The initial data is a rational function $R(x) := B(x)/A(x)$. We assume that A and B are polynomials with real coefficients and A has no real zeros and write

$$(5.4) \quad A(x) := \sum_{k=0}^p a_k x^{p-k} \text{ and } B(x) := \sum_{k=0}^{p-2} b_k x^{p-2-k}.$$

Step 2. Choose a positive integer $m \geq 2$.

Step 3. Introduce the polynomial

$$(5.5) \quad H(x) := \text{Res}_z(A(z), P_m(z) - xQ_m(z))$$

and write it as

$$(5.6) \quad H(x) := \sum_{l=0}^p e_l x^{p-l}.$$

The polynomial H is thus defined as the determinant of the Sylvester matrix which is formed of the polynomial coefficients. As such, the coefficients e_l of $H(x)$ themselves are integer polynomials in the a_i . Explicitly,

$$(5.7) \quad e_l = (-1)^l a_0^m \prod_{j=1}^p Q_m(x_j) \times \sigma_l^{(p)}(R_m(x_1), R_m(x_2), \dots, R_m(x_p)),$$

where $\{x_1, x_2, \dots, x_p\}$ are the roots of A , each written according to multiplicity. The functions $\sigma_l^{(p)}$ are the elementary symmetric functions in p variables defined by

$$(5.8) \quad \prod_{l=1}^p (y - y_l) = \sum_{l=0}^p (-1)^l \sigma_l^{(p)}(y_1, \dots, y_p) y^{p-l}.$$

It is possible to compute the coefficients e_l symbolically from the coefficients of A , without the knowledge of the roots of A .

Also define

$$(5.9) \quad E(x) := H(R_m(x)) \times Q_m(x)^p.$$

Step 4. The polynomial A divides E and we denote the quotient by Z . The coefficients of Z are integer polynomials in the a_i .

Step 5. Define the polynomial $C(x) := B(x)Z(x)$.

Step 6. There exists a polynomial $J(x)$, whose coefficients have an explicit formula in terms of the coefficients c_j of $C(x)$, such that

$$(5.10) \quad \int_{-\infty}^{\infty} \frac{B(x)}{A(x)} dx = \int_{-\infty}^{\infty} \frac{J(x)}{H(x)} dx.$$

This new integrand is the rational function whose existence is guaranteed by Lemma 4.1. The explicit computation of the coefficients of J can be found in [41]. This is the *rational Landen transformation* of order m .

Example 5.1. Completing the algorithm with $m = 3$ and the rational function

$$(5.11) \quad R(x) = \frac{1}{ax^2 + bx + c},$$

produces the result stated below. Notice that the values of the iterates are ratios of integer polynomials of degree 3, as was stated above. The details of this example appear in [42].

Theorem 5.2. The integral

$$(5.12) \quad I = \int_{-\infty}^{\infty} \frac{dx}{ax^2 + bx + c}$$

is invariant under the transformation

$$(5.13) \quad \begin{aligned} a &\mapsto \frac{a}{\Delta} ((a+3c)^2 - 3b^2), \\ b &\mapsto \frac{b}{\Delta} (3(a-c)^2 - b^2), \\ c &\mapsto \frac{c}{\Delta} ((3a+c)^2 - 3b^2), \end{aligned}$$

where $\Delta := (3a+c)(a+3c) - b^2$. The condition $b^2 - 4ac < 0$, imposed to ensure convergence of the integral, is preserved by the iteration.

Example 5.2. In this example we follow the steps described above in order to produce a rational Landen transformation of order 2 for the integral

$$(5.14) \quad I = \int_{-\infty}^{\infty} \frac{b_0x^4 + b_1x^3 + b_2x^2 + b_3x + b_4}{a_0x^6 + a_1x^5 + a_2x^4 + a_3x^3 + a_4x^2 + a_5x + a_6} dx.$$

Recall that the algorithm starts with a rational function $R(x)$ and produces a new function $\mathfrak{L}_2(R(x))$ satisfying

$$(5.15) \quad \int_{-\infty}^{\infty} R(x) dx = \int_{-\infty}^{\infty} \mathfrak{L}_2(R(x)) dx.$$

Step 1. The initial data is $R(x) = B(x)/A(x)$ with

$$(5.16) \quad A(x) = a_0x^6 + a_1x^5 + a_2x^4 + a_3x^3 + a_4x^2 + a_5x + a_6,$$

and

$$(5.17) \quad B(x) = b_0x^4 + b_1x^3 + b_2x^2 + b_3x + b_4.$$

The parameter p is the degree of A , so $p = 6$.

Step 2. We choose $m = 2$ to produce a method of order 2. The algorithm employs the polynomials $P_2(z) = z^2 - 1$ and $Q_2(z) = 2z$.

Step 3. The polynomial

$$(5.18) \quad H(x) := \text{Res}_z(A(z), z^2 - 1 - 2xz)$$

is computed with the Mathematica command `Resultant` to obtain

$$(5.19) \quad H(x) = e_0x^6 + e_1x^5 + e_2x^4 + e_3x^3 + e_4x^2 + e_5x + e_6,$$

with

$$\begin{aligned}
 (5.20) \quad e_0 &= 64a_0a_6, \\
 e_1 &= -32(a_0a_5 - a_1a_6), \\
 e_2 &= 16(a_0a_4 - a_1a_5 + 6a_0a_6 + a_2a_6), \\
 e_3 &= -8(a_0a_3 - a_1a_4 + 5a_0a_5 + a_2a_5 - 5a_1a_6 - a_3a_6), \\
 e_4 &= 4(a_0a_2 - a_1a_3 + 4a_0a_4 + a_2a_4 - 4a_1a_5 - a_3a_5 + 9a_0a_6 + 4a_2a_6 + a_4a_6), \\
 e_5 &= -2(a_0a_1 - a_1a_2 + 3a_0a_3 + a_2a_3 - 3a_1a_4 - a_3a_4 + 5a_0a_5 + \\
 &\quad + 3a_2a_5 + a_4a_5 - 5a_1a_6 - 3a_3a_6 - a_5a_6), \\
 e_6 &= (a_0 - a_1 + a_2 - a_3 + a_4 - a_5 + a_6)(a_0 + a_1 + a_2 + a_3 + a_4 + a_5 + a_6).
 \end{aligned}$$

The polynomial $H(x)$ is the denominator of the integrand $\mathfrak{L}_2(R(x))$ in (5.15).

In Step 3 we also define

$$(5.21) \quad E(x) = H(R_2(x))Q_2^6(x) = H\left(\frac{x^2 - 1}{2x}\right) \cdot (2x)^6.$$

The function $E(x)$ is a polynomial of degree 12, written as

$$(5.22) \quad E(x) = \sum_{k=0}^{12} \alpha_k x^{12-k}.$$

Using the expressions for e_j in (5.20) in (5.21) yields

$$\begin{aligned}
 (5.23) \quad \alpha_0 &= \alpha_{12} = 64a_0a_6 \\
 \alpha_1 &= -\alpha_{11} = -64(a_0a_5 - a_1a_6), \\
 \alpha_2 &= \alpha_{10} = 64(a_0a_4 - a_1a_5 + a_2a_6), \\
 \alpha_3 &= -\alpha_9 = -64(a_0a_3 - a_1a_4 + a_2a_5 - a_3a_6), \\
 \alpha_4 &= \alpha_8 = 64(a_0a_2 - a_1a_3 + a_2a_4 - a_3a_5 + a_4a_6), \\
 \alpha_5 &= -\alpha_7 = -64(a_0a_1 - a_1a_2 + a_2a_3 - a_3a_4 + a_4a_5 - a_5a_6), \\
 \alpha_6 &= 64(a_0^2 - a_1^2 + a_2^2 - a_3^2 + a_4^2 - a_5^2 + a_6^2).
 \end{aligned}$$

Step 4. The polynomial $A(x)$ always divides $E(x)$. The quotient is denoted by $Z(x)$. The values of α_j given in (5.23) produce

$$(5.24) \quad Z(x) = 64(a_0 - a_1x + a_2x^2 - a_3x^3 + a_4x^4 - a_5x^5 + a_6x^6).$$

Step 5. Define the polynomial $C(x) := B(x)Z(x)$. In this case, C is of degree 10, written as

$$(5.25) \quad C(x) = \sum_{k=0}^{10} c_k x^{10-k},$$

and the coefficients c_k are given by

(5.26)

$$\begin{aligned}
c_0 &= 64a_6b_0, \\
c_1 &= -64(a_5b_0 - a_6b_1), \\
c_2 &= 64(a_4b_0 - a_5b_1 + a_6b_2), \\
c_3 &= -64(a_3b_0 - a_4b_1 + a_5b_2 - a_6b_3), \\
c_4 &= 64(a_2b_0 - a_3b_1 + a_4b_2 - a_5b_3 + a_6b_4), \\
c_5 &= -64(a_1b_0 - a_2b_1 + a_3b_2 - a_4b_3 + a_5b_4), \\
c_6 &= 64(a_0b_0 - a_1b_1 + a_2b_2 - a_3b_3 + a_4b_4), \\
c_7 &= 64(a_0b_1 - a_1b_2 + a_2b_3 - a_3b_4), \\
c_8 &= 64(a_0b_2 - a_1b_3 + a_2b_4), \\
c_9 &= 64(a_0b_3 - a_1b_4), \\
c_{10} &= 64a_0b_4.
\end{aligned}$$

Step 6 produces the numerator $J(x)$ of the new integrand $\mathfrak{L}_2(R(x))$ from the coefficients c_j given in (5.26). The function $J(x)$ is a polynomial of degree 4, written as

$$(5.27) \quad J(x) = \sum_{k=0}^4 j_k x^{4-k}.$$

Using the values of (5.26) we obtain

(5.28)

$$\begin{aligned}
j_0 &= 32(a_6b_0 + a_0b_4), \\
j_1 &= -16(a_5b_0 - a_6b_1 + a_0b_3 - a_1b_4), \\
j_2 &= 8(a_4b_0 + 3a_6b_0 - a_5b_1 + a_0b_2 + a_6b_2 - a_1b_3 + 3a_0b_4 + a_2b_4), \\
j_3 &= -4(a_3b_0 + 2a_5b_0 + a_0b_1 - a_4b_1 - 2a_6b_1 - a_1b_2 + a_5b_2 + \\
&\quad + 2a_0b_3 + a_3b_3 - a_6b_3 - 2a_1b_4 - a_3b_4), \\
j_4 &= 2(a_0b_0 + a_2b_0 + a_4b_0 + a_6b_0 - a_1b_1 - a_3b_1 - a_5b_1 + a_0b_2 + \\
&\quad + a_2b_2 + a_4b_2 + a_6b_2 - a_1b_3 - a_3b_3 - a_5b_3 + a_0b_4 + a_2b_4 + a_4b_4 + a_6b_4).
\end{aligned}$$

The explicit computation of the coefficients of J can be found in [41].

The new rational function is

$$(5.29) \quad \mathfrak{L}_2(R(x)) := \frac{J(x)}{H(x)},$$

with $J(x)$ given in (5.27) and $H(x)$ in (5.19). The transformation is

$$\frac{b_0x^4 + b_1x^3 + b_2x^2 + b_3x + b_4}{a_0x^6 + a_1x^5 + a_2x^4 + a_3x^3 + a_4x^2 + a_5x + a_6} \mapsto \frac{j_0x^4 + j_1x^3 + j_2x^2 + j_3x + j_4}{e_0x^6 + e_1x^5 + e_2x^4 + e_3x^3 + e_4x^2 + e_5x + e_6}.$$

The numerator coefficients are given in (5.20) and the denominator ones in (5.28), explicitly as polynomials in the coefficients of the original rational function. The generation of these polynomials is a completely symbolic procedure.

The first two steps of this algorithm, applied to the definite integral

$$(5.30) \quad \int_{-\infty}^{\infty} \frac{dx}{x^6 + x^3 + 1} = \frac{\pi}{9} \left(2\sqrt{3} \cos(\pi/9) + \sqrt{3} \cos(2\pi/9) + 3 \sin(2\pi/9) \right),$$

produces the identities

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{dx}{x^6 + x^3 + 1} &= \int_{-\infty}^{\infty} \frac{2(16x^4 + 12x^2 + 2x + 2)}{64x^6 + 96x^4 + 36x^2 + 3} dx \\ &= \int_{-\infty}^{\infty} \frac{4(2816x^4 - 1024x^3 + 8400x^2 - 884x + 5970)}{12288x^6 + 59904x^4 + 87216x^2 + 39601} dx. \end{aligned}$$

The convergence of the iterations of rational Landen transformations is discussed in the next section.

6. THE ISSUE OF CONVERGENCE

The convergence of the double sequence (a_n, b_n) appearing in the elliptic Landen transformation (1.1) is easily established. Assume $0 < b_0 \leq a_0$, then the arithmetic-geometric inequality yields $b_n \leq b_{n+1} \leq a_{n+1} \leq a_n$. Also

$$(6.1) \quad 0 \leq a_{n+1} - b_{n+1} = \frac{1}{2} \frac{(a_n - b_n)^2}{(\sqrt{a_n} + \sqrt{b_n})^2}.$$

This shows that a_n and b_n have a common limit: $M = \text{AGM}(a, b)$, the arithmetic-geometric of a and b . The convergence is quadratic:

$$(6.2) \quad |a_{n+1} - M| \leq C |a_n - M|^2,$$

for some constant $C > 0$ independent of n . Details can be found in [15].

The Landen transformations produce maps on the space of coefficients of the integrand. In this section, we discuss the convergence of the rational Landen transformations. This discussion is divided in two cases:

Case 1: the half-line. Let $R(x)$ be an even rational function, written as $R(x) = P(x)/Q(x)$, with

$$(6.3) \quad P(x) = \sum_{k=0}^{p-1} b_k x^{2(p-1-k)}, \quad Q(x) = \sum_{k=0}^p a_k x^{2(p-k)},$$

and $a_0 = a_p = 1$. The *parameter space* is

$$(6.4) \quad \mathfrak{P}_{2p}^+ = \{(a_1, \dots, a_{p-1}; b_0, \dots, b_{p-1})\} \subset \mathbb{R}^{p-1} \times \mathbb{R}^p.$$

We write

$$(6.5) \quad \mathbf{a} := (a_1, \dots, a_{p-1}) \text{ and } \mathbf{b} := (b_0, \dots, b_p).$$

Define

$$(6.6) \quad \Lambda_{2p} = \{(a_1, \dots, a_{p-1}) \in \mathbb{R}^{p-1} : \int_0^\infty R(x) dx \text{ is finite}\}.$$

Observe that the convergence of the integral depends only on the parameters in the denominator.

The Landen transformations provide a map

$$(6.7) \quad \Phi_{2p} : \mathfrak{P}_{2p}^+ \rightarrow \mathfrak{P}_{2p}^+$$

that preserves the integral. Introduce the notation

$$(6.8) \quad \mathbf{a}_n = (a_1^{(n)}, \dots, a_{p-1}^{(n)}) \text{ and } \mathbf{b}_n = (b_0^{(n)}, \dots, b_p^{(n)}),$$

where

$$(6.9) \quad (\mathbf{a}_n, \mathbf{b}_n) = \Phi_{2p}(\mathbf{a}_{n-1}, \mathbf{b}_{n-1})$$

are the iterates of the map Φ_{2p} .

The result that one expects is this:

Theorem 6.1. The region Λ_{2p} is invariant under the map Φ_{2p} . Moreover

$$(6.10) \quad \mathbf{a}_n \rightarrow \left(\binom{p}{1}, \binom{p}{2}, \dots, \binom{p}{p-1} \right),$$

and there exists a number L , that depends on the initial conditions, such that

$$(6.11) \quad \mathbf{b}_n \rightarrow \left(\binom{p-1}{0} L, \binom{p-1}{1} L, \dots, \binom{p-1}{p-1} L \right).$$

This is equivalent to say that the sequence of rational functions formed by the Landen transformations, converge to $L/(x^2 + 1)$.

This was established in [33] using the geometric language of Landen transformations which, while unexpected, is satisfactory.

Theorem 6.2. Let φ be a 1-form, holomorphic in a neighborhood of $\mathbb{R} \subset \mathbb{P}^1$. Then

$$(6.12) \quad \lim_{n \rightarrow \infty} (\pi_*)^n \varphi = \frac{1}{\pi} \left(\int_{-\infty}^{\infty} \varphi \right) \frac{dz}{1+z^2},$$

where the convergence is uniform on compact subsets of U , the neighborhood in the definition of π_* .

The proof is detailed for the map $\pi(z) = \frac{z^2-1}{2z} = R_2(z)$, but it extends without difficulty to the generalization R_m .

Theorem 6.2 can be equivalently reformulated as:


Theorem 6.3. The iterates of the Landen transformation starting at $(\mathbf{a}_0, \mathbf{b}_0) \in \mathfrak{P}_{2p}^+$ converge (to the limit stated in Theorem 6.1) if and only if the integral formed by the initial data is finite.

It would be desirable to establish this result by purely dynamical techniques. This has been established only for the case $p = 3$. In that case the Landen transformation for

$$(6.13) \quad U_6 := \int_0^\infty \frac{cx^4 + dx^2 + e}{x^6 + ax^4 + bx^2 + 1} dx$$

is

$$(6.14) \quad \begin{aligned} a_1 &\leftarrow \frac{ab + 5a + 5b + 9}{(a + b + 2)^{4/3}}, \\ b_1 &\leftarrow \frac{a + b + 6}{(a + b + 2)^{2/3}}, \end{aligned}$$

FIGURE 1. Zero locus of $R(a, b)$

coupled with

$$\begin{aligned} c_1 &\leftarrow \frac{c + d + e}{(a + b + 2)^{2/3}}, \\ d_1 &\leftarrow \frac{(b + 3)c + 2d + (a + 3)e}{a + b + 2}, \\ e_1 &\leftarrow \frac{c + e}{(a + b + 2)^{1/3}}. \end{aligned}$$

The region

$$(6.15) \quad \Lambda_6 = \{(a, b) \in \mathbb{R}^2 : U_6 < \infty\}$$

is described by the discriminant curve $\mathfrak{R}$, the zero set of the polynomial

$$(6.16) \quad R(a, b) = 4a^3 + 4b^3 - 18ab - a^2b^2 + 27.$$

This zero set, shown in Figure 1, has two connected components: the first one $\mathfrak{R}_+$ contains $(3, 3)$ as a cusp and the second one $\mathfrak{R}_-$, given by $R_-(a, b) = 0$, is disjoint from the first quadrant. The branch $\mathfrak{R}_-$ is the boundary of the set Λ_6 .

The identity

$$(6.17) \quad R(a_1, b_1) = \frac{(a - b)^2 R(a, b)}{(a + b + 2)^4},$$

shows that $\partial\mathfrak{R}$ is invariant under Φ_6 . By examining the effect of this map along lines of slope -1 , we obtain a direct parametrization of the flow *on* the discriminant curve. Indeed, this curve is parametrized by

$$(6.18) \quad a(s) = \frac{s^3 + 4}{s^2} \text{ and } b(s) = \frac{s^3 + 16}{4s}.$$

Then

$$(6.19) \quad \varphi(s) = \left(\frac{4(s^2 + 4)^2}{s(s+2)^2} \right)^{1/3}$$

gives the image of the Landen transformation Φ_6 ; that is,

$$(6.20) \quad \Phi_6(a(s), b(s)) = (a(\varphi(s)), b(\varphi(s))).$$

The map Φ_6 has three fixed points: $(3, 3)$, that is super-attracting, a saddle point P_2 on the lower branch $\mathfrak{R}_-$ of the discriminant curve, and a third unstable spiral below this lower branch. In [25] we prove:

Theorem 6.4. The lower branch of the discriminant curve is the curve Λ_6 . This curve is also the global unstable manifold of the saddle point P_2 . Therefore the iterations of Φ_6 starting at (a, b) converge if and only if the integral U_6 , formed with the parameters (a, b) , is finite. Moreover, $(a_n, b_n) \rightarrow (3, 3)$ quadratically and there exists a number L such that $(c_n, d_n, e_n) \rightarrow (1, 2, 1)L$.

The next result provides an analogue of the AGM (1.12) for the rational case. The main differences here are that our iterates converge to an *algebraic* number and we achieve *order- m* convergence.

Case 2: The whole-line: This works for any choice of positive integer m . Let $R(x)$ be a rational function, written as $R(x) = B(x)/A(x)$. Assume that the coefficients of A and B are real, that A has no real zeros and that $\deg(B) \leq \deg(A) - 2$. These conditions are imposed to guarantee the existence of

$$(6.21) \quad I = \int_{-\infty}^{\infty} R(x) dx.$$

In particular A must be of even degree, and we write

$$(6.22) \quad A(x) = \sum_{k=0}^p a_k x^{p-k} \text{ and } B(x) = \sum_{k=0}^{p-2} b_k x^{p-2-k}.$$

We can also require that $\deg(\gcd(A, B)) = 0$.

The class of such rational functions will be denoted by $\mathfrak{R}_p$.

The algorithm presented in Section 5 provides a transformation on the parameters

$$(6.23) \quad \mathfrak{P}_p := \{a_0, a_1, \dots, a_p; b_0, b_1, \dots, b_{p-2}\} = \mathbb{R}^{p+1} \times \mathbb{R}^{p-1}$$

of $R \in \mathfrak{R}_p$ that preserves the integral I . In fact, we produce a family of maps, indexed by $m \in \mathbb{N}$,

$$\mathfrak{L}_{m,p} : \mathfrak{R}_p \rightarrow \mathfrak{R}_p,$$

such that

$$(6.24) \quad \int_{-\infty}^{\infty} R(x) dx = \int_{-\infty}^{\infty} \mathfrak{L}_{m,p}(R(x)) dx.$$

The maps $\mathfrak{L}_{m,p}$ induce a *rational Landen transformation* on the parameter space:

$$(6.25) \quad \Phi_{m,p} : \mathfrak{P}_p \rightarrow \mathfrak{P}_p$$

by simply listing the coefficients of the function $\mathfrak{L}_{m,p}(R(x))$.

The original integral is written in the form

$$(6.26) \quad I = \frac{b_0}{a_0} \int_{-\infty}^{\infty} \frac{x^{p-2} + b_0^{-1}b_1x^{p-3} + b_0^{-1}b_2x^{p-4} + \cdots + b_0^{-1}b_{p-2}}{x^p + a_0^{-1}a_1x^{p-1} + a_0^{-1}a_2x^{p-2} + \cdots + a_0^{-1}a_p} dx.$$

The Landen transformation generates a sequence of coefficients,

$$(6.27) \quad \mathfrak{P}_{p,n} := \{a_0^{(n)}, a_1^{(n)}, \dots, a_p^{(n)}; b_0^{(n)}, b_1^{(n)}, \dots, b_{p-2}^{(n)}\},$$

with $\mathfrak{P}_{p,0} = \mathfrak{P}_p$ as in (6.23). We expect that, as $n \rightarrow \infty$,

$$(6.28) \quad \mathbf{x}_n := \left(\frac{a_1^{(n)}}{a_0^{(n)}}, \frac{a_2^{(n)}}{a_0^{(n)}}, \dots, \frac{a_p^{(n)}}{a_0^{(n)}}, \frac{b_1^{(n)}}{b_0^{(n)}}, \frac{b_2^{(n)}}{b_0^{(n)}}, \dots, \frac{b_{p-2}^{(n)}}{b_0^{(n)}} \right)$$

converges to

$$(6.29) \quad \mathbf{x}_\infty := \left(0, \binom{q}{1}, 0, \binom{q}{2}, \dots, \binom{q}{q}; 0, \binom{q-1}{1}, 0, \binom{q-1}{2}, \dots, \binom{q-1}{q-1} \right),$$

where $q = p/2$. Moreover, we should have

$$(6.30) \quad \|\mathbf{x}_{n+1} - \mathbf{x}_\infty\| \leq C \|\mathbf{x}_n - \mathbf{x}_\infty\|^m.$$

The invariance of the integral then shows that

$$(6.31) \quad \frac{b_0^{(n)}}{a_0^{(n)}} \rightarrow \frac{1}{\pi} I.$$

This produces an iterative method to evaluate the integral of a rational function. The method's convergence is of order- m .

The convergence of these iterations, and in particular the bound (6.30), can be established by the argument presented in section 4. Thus, the transformation $\mathfrak{L}_{m,p}$ leads to a sequence that has order- m convergence. We expect to develop these ideas into an efficient numerical method for integration.

We choose to measure the convergence of a sequence of vectors to 0 is in the L_2 -norm,

$$(6.32) \quad \|v\|_2 = \frac{1}{\sqrt{2p-2}} \left(\sum_{k=1}^{2p-2} \|v_k\|^2 \right)^{1/2},$$

and also the L_∞ -norm,

$$(6.33) \quad \|v\|_\infty = \text{Max} \{ \|v_k\| : 1 \leq k \leq 2p-2 \}.$$

The rational functions appearing as integrands have rational coefficients, so, as a measure of their complexity, we take the largest number of digits of these coefficients. This appears in the column marked *size*.

The following tables illustrate the iterates of rational Landen transformations of order 2, 3 and 4, applied to the example

$$I = \int_{-\infty}^{\infty} \frac{3x+5}{x^4+14x^3+74x^2+184x+208} dx = -\frac{7\pi}{12}.$$

The first column gives the L_2 -norm of $u_n - u_\infty$, the second its L_∞ -norm, the third presents the relative error in (6.31), and in the last column we give the size of the rational integrand. At each step, we verify that the new rational function integrates to $-7\pi/12$.

Method of order 2

n	L_2 -norm	L_∞ -norm	Error	Size
1	58.7171	69.1000	1.02060	5
2	7.444927	9.64324	1.04473	10
3	4.04691	5.36256	0.945481	18
4	1.81592	2.41858	1.15092	41
5	0.360422	0.411437	0.262511	82
6	0.0298892	0.0249128	0.0189903	164
7	0.000256824	0.000299728	0.0000362352	327
8	1.92454×10^{-8}	2.24568×10^{-8}	1.47053×10^{-8}	659
9	1.0823×10^{-16}	1.2609×10^{-16}	8.2207×10^{-17}	1318

As expected, for the method of order 2, we observe quadratic convergence in the L_2 -norm and also in the L_∞ -norm. The size of the coefficients of the integrand is approximately doubled at each iteration.

Method of order 3

n	L_2 -norm	L_∞ -norm	Error	Size
1	15.2207	20.2945	1.03511	8
2	1.97988	1.83067	0.859941	23
3	0.41100	0.338358	0.197044	69
4	0.00842346	0.00815475	0.00597363	208
5	5.05016×10^{-8}	5.75969×10^{-8}	1.64059×10^{-9}	626
6	1.09651×10^{-23}	1.02510×10^{-23}	3.86286×10^{-24}	1878
7	1.12238×10^{-70}	1.22843×10^{-70}	8.59237×10^{-71}	5634

Method of order 4

n	L_2 -norm	L_∞ -norm	Error	Size
1	7.44927	9.64324	1.04473	10
2	1.81592	2.41858	1.15092	41
3	0.0298892	0.0249128	0.0189903	164
4	1.92454×10^{-8}	2.249128×10^{-8}	1.47053×10^{-8}	659
5	3.40769×10^{-33}	3.96407×10^{-33}	2.56817×10^{-33}	2637

Example 6.1. A method of order 3 for the evaluation of the quadratic integral

$$(6.34) \quad I = \int_{-\infty}^{\infty} \frac{dx}{ax^2 + bx + c},$$

has been analyzed in [42]. We refer to Example 5.1 for the explicit formulas of this Landen transformation, and define the iterates accordingly. From there, we prove that the error term,

$$(6.35) \quad e_n := \left(a_n - \frac{1}{2}\sqrt{4ac - b^2}, b_n, c_n - \frac{1}{2}\sqrt{4ac - b^2} \right)$$

satisfies $e_n \rightarrow 0$ as $n \rightarrow \infty$, with cubic rate:

$$(6.36) \quad \|e_{n+1}\| \leq C\|e_n\|^3.$$

The proof of convergence is elementary. Therefore we have

$$(6.37) \quad (a_n, b_n, c_n) \rightarrow (\sqrt{ac - b^2/4}, 0, \sqrt{4ac - b^2/4}) ,$$

which, in conjunction with (6.34), implies that

$$(6.38) \quad I = \frac{2}{\sqrt{4ac - b^2}} \int_{-\infty}^{\infty} \frac{dx}{x^2 + 1} ,$$

exactly as one would have concluded by completing the square. Unlike completing the square, our method extends to a general rational integral over the real line.

7. THE APPEARANCE OF THE AGM IN DIVERSE CONTEXTS

The (elliptic) Landen transformation

$$(7.1) \quad a_1 \leftarrow \frac{1}{2}(a + b) \text{ and } b_1 \leftarrow \sqrt{ab}$$

leaving invariant the elliptic integral

$$(7.2) \quad G(a, b) = \int_0^{\pi/2} \frac{d\varphi}{\sqrt{a^2 \cos^2 \varphi + b^2 \sin^2 \varphi}}$$

appears in many different forms. In this last section we present a partial list of them.

7.1. The Elliptic Landen Transformation. For the lattice $\mathbb{L} = \mathbb{Z} \oplus \omega\mathbb{Z}$, introduce the *theta-functions*

$$(7.3) \quad \vartheta_3(x, \omega) := \sum_{n=-\infty}^{\infty} z^{2n} q^{n^2} \text{ and } \vartheta_4(x, \omega) := \sum_{n=-\infty}^{\infty} (-1)^n z^{2n} q^{n^2},$$

where $z = e^{\pi i x}$ and $q = e^{\pi i \omega}$. The condition $\text{Im } \omega > 0$ is imposed to ensure convergence of the series. These functions admit a variety of remarkable identities. In particular, the *null-values* (those with $x = 0$) satisfy

$$\vartheta_4^2(0, 2\omega) = \vartheta_3(0, \omega)\vartheta_4(0, \omega) \text{ and } \vartheta_3^2(0, 2\omega) = \frac{1}{2} (\vartheta_3^2(0, \omega) + \vartheta_4^2(0, \omega)) ,$$

and completely characterize values of the AGM, leading to the earlier result [15]. Grayson [30] has used the doubling of the period ω to derive the arithmetic-geometric mean from the cubic equations describing the corresponding elliptic curves. See Chapter 3 in [43] for more information. P. Sole et al. [49, 50] have produced proofs of generalizations of these identities by lattice enumeration methods related to binary and ternary codes.

7.2. A time-one map. We now present a deeper and more modern version of a result known to Gauss: given a sequence of points $\{x_n\}$ on a manifold X , decide whether there is a differential equation

$$(7.4) \quad \frac{dx}{dt} = V(x),$$

starting at x_0 such that $x_n = x(n, x_0)$. Here $x(t, x_0)$ is the unique solution to (7.4) satisfying $x(0, x_0) = x_0$. Denote by

$$(7.5) \quad \phi_{\text{ellip}}(a, b) = \left(\frac{1}{2}(a+b), \sqrt{ab} \right)$$

the familiar elliptic Landen transformation. Now take $a, b \in \mathbb{R}$ with $a > b > 0$. Use the null-values of the theta functions to find unique values (τ, ρ) such that

$$(7.6) \quad a = \rho \vartheta_3^2(0, \tau) \text{ and } b = \rho \vartheta_4^2(0, \tau).$$

Finally define

$$(7.7) \quad x_{\text{ellip}}(t) = (a(t), b(t)) = \rho \left(\vartheta_3^2(0, 2^t \tau), \vartheta_4^2(0, 2^t \tau) \right),$$

with $x_{\text{ellip}}(0) = (a, b)$. The remarkable result is [28]:

Theorem 7.1. (Deift, Li, Previato, Tomei). The map $t \rightarrow x_{\text{ellip}}(t)$ is an integrable Hamiltonian flow on X equipped with an appropriate symplectic structure. The Hamiltonian is the complete elliptic integral $G(a, b)$ and the angle is (essentially the logarithm of) the second period of the elliptic curve associated with τ . Moreover

$$(7.8) \quad x_{\text{ellip}}(k) = \phi_{\text{ellip}}^k(a, b).$$

Thus the arithmetic-geometric algorithm is the time-one map of a completely integrable Hamiltonian flow.

Notice that this theorem shows that the result in question respects some additional structures whose invention postdates Gauss.

A natural question is whether the map (3.9) appears as a time-one map of an interesting flow.

7.3. A quadruple sequence. Several variations of the elliptic Landen appear in the literature. Borchardt [5] considers the four-term quadratically convergent iteration

$$(7.9) \quad \begin{aligned} a_{n+1} &= \frac{a_n + b_n + c_n + d_n}{4}, \\ b_{n+1} &= \frac{\sqrt{a_n b_n} + \sqrt{c_n d_n}}{2}, \\ c_{n+1} &= \frac{\sqrt{a_n c_n} + \sqrt{b_n d_n}}{2}, \\ d_{n+1} &= \frac{\sqrt{a_n d_n} + \sqrt{b_n c_n}}{2}, \end{aligned}$$

starting with $a_0 = a$, $b_0 = b$, $c_0 = c$ and $d_0 = d$. The common limit, denoted by $G(a, b, c, d)$, is given by

$$(7.10) \quad \frac{1}{G(a, b, c, d)} = \frac{1}{\pi^2} \int_0^{\alpha_3} \int_{\alpha_1}^{\alpha_2} \frac{(x-y) dx dy}{\sqrt{R(x)R(y)}},$$

where $R(x) = x(x - \alpha_0)(x - \alpha_1)(x - \alpha_2)(x - \alpha_3)$ and the numbers α_j are given by explicit formulas in terms of the parameters a, b, c, d . Details are given by Mestre in [44].

The initial conditions $(a, b, c, d) \in \mathbb{R}^4$ for which the iteration converges has some interesting invariant subsets. When $a = b$ and $c = d$, we recover the AGM iteration (1.1). In the case that $b = c = d$, we have another invariant subset, linking to an iterative mean described below.

7.4. Variations of AGM with hypergeometric limit. Let $N \in \mathbb{N}$. The analysis of

$$(7.11) \quad a_{n+1} = \frac{a_n + (N-1)b_n}{N} \text{ and } c_{n+1} = \frac{a_n - b_n}{N},$$

with $b_n = (a_n^N - c_n^N)^{1/N}$, is presented in [18]. All the common ingredients appear there: a common limit, fast convergence, theta functions and sophisticated iterations for the evaluation of π . The common limit is denoted by $\text{AG}_N(a, b)$. The convergence is of order N and the limit is identified for small N : for $0 < k < 1$,

$$\begin{aligned} \frac{1}{\text{AG}_2(1, k)} &= {}_2F_1(1/2, 1/2; 1; 1 - k^2) \text{ and} \\ \frac{1}{\text{AG}_3(1, k)} &= {}_2F_1(1/3, 2/3; 1; 1 - k^2). \end{aligned}$$

where

$$(7.12) \quad {}_2F_1(a, b; c; x) = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(c)_k k!} x^k$$

is the classical hypergeometric function. There are integral representations of these as well which parallel (1.12), see [13], Section 6.1 for details.

Other hypergeometric values appear from similar iterations. For example,

$$(7.13) \quad a_{n+1} = \frac{a_n + 3b_n}{4} \text{ and } b_{n+1} = \sqrt{b_n(a_n + b_n)/2},$$

have a common limit, denoted by $A_4(a, b)$. It is given by

$$(7.14) \quad \frac{1}{A_4(1, k)} = {}_2F_1^2(1/4, 3/4; 1; 1 - k^2).$$

To compute π quartically, start at $a_0 = 1, b_0 = (12\sqrt{2} - 16)^{1/4}$. Now compute a_n from two steps of AG_2 :

$$(7.15) \quad a_{n+1} = \frac{a_n + b_n}{2}, \text{ and } b_{n+1} = \left(\frac{a_n b_n^3 + b_n a_n^3}{2} \right)^{1/4}.$$

Then

$$(7.16) \quad \pi = \lim_{n \rightarrow \infty} 3a_{n+1}^4 \left(1 - \sum_{j=0}^n 2^{j+1} (a_j^4 - a_{j+1}^4) \right)^{-1}$$

with $|a_{n+1} - \pi| \leq C|a_n - \pi|^4$, for some constant $C > 0$. This is much better than the partial sums of

$$(7.17) \quad \pi = 4 \sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1}.$$

The sequence (a_n, b_n) defined by the iteration

$$(7.18) \quad a_{n+1} = \frac{a_n + 2b_n}{3}, \quad b_{n+1} = \left(\frac{b_n(a_n^2 + a_nb_n + b_n^2)}{3} \right)^{\frac{1}{3}},$$

starting at $a_0 = 1$, $b_0 = x$ are analyzed in [17]. They have a common limit $F(x)$ given by

$$(7.19) \quad \frac{1}{F(x)} = {}_2F_1\left(\frac{1}{3}, \frac{2}{3}; 1; 1 - x^3\right).$$

7.5. Iterations where the limit is harder to find. J. Borwein and P. Borwein [16] studied the iteration of

$$(7.20) \quad (a, b) \rightarrow \left(\frac{a + 3b}{4}, \frac{\sqrt{ab} + b}{2} \right),$$

and showed the existence of a common limit $B(a_0, b_0)$. Define $B(x) = B(1, x)$. The study of the iteration (7.20) is based on the functional equation

$$(7.21) \quad B(x) = \frac{1 + 3x}{4} B\left(\frac{2(\sqrt{x} + x)}{1 + 3x}\right).$$

and a parametrization of the iterates by theta functions [16]. The complete analysis of (7.20) starts with the purely computational observation that

$$(7.22) \quad B(x) \sim \frac{\pi^2}{3} \log^{-2}(x/4) \text{ as } x \rightarrow 0.$$

H. H. Chan, K. Chua and P. Sole [31] identified the limiting function as

$$(7.23) \quad B(x) = \left({}_2F_1\left(\frac{1}{3}, \frac{1}{6}; 1; 27 \frac{x(1-x)^2}{(1+3x)^3}\right) \right)^{-2},$$

valid for $\frac{2}{3} < x < 1$. A similar hypergeometric expression gives $B(x)$ for $0 < x < \frac{2}{3}$.

7.6. Fast computation of elementary functions. The fast convergence of the elliptic Landen recurrence (1.1) to the arithmetic-geometric mean provides a method for numerical evaluation of the elliptic integral $G(a, b)$. The same idea provides for the fast computation of elementary functions. For example, in [14] we find the estimate

$$(7.24) \quad \left| \log x - (G(1, 10^{-n}) - G(1, 10^{-n}x)) \right| < n10^{-2(n-1)},$$

for $0 < x < 1$ and $n \geq 3$.

7.7. A continued fraction. The continued fraction

$$(7.25) \quad R_\eta(a, b) = \frac{a}{b^2 + \frac{\eta + \frac{4a^2}{\eta + \frac{9b^2}{\eta + \dots}}}{\eta + \frac{4a^2}{\eta + \frac{9b^2}{\eta + \dots}}}},$$

has an interesting connection to the AGM. In their study of the convergence of $R_\eta(a, b)$, J. Borwein, R. Crandall and G. Fee [19] established the identity

$$(7.26) \quad R_\eta\left(\frac{a+b}{2}, \sqrt{ab}\right) = \frac{1}{2} (R_\eta(a, b) + R_\eta(b, a)).$$

This identity originates with Ramanujan; the similarity with AGM is now direct.

The continued fraction converges for positive real parameters, but for $a, b \in \mathbb{C}$ the convergence question is quite delicate. For example, the even/odd parts of $R_1(1, i)$ converge to distinct limits. See [19] and [20] for more details.

7.8. Elliptic Landen with complex initial conditions. The iteration of (1.1) with $a_0, b_0 \in \mathbb{C}$ requires a choice of square root at each step. Let $a, b \in \mathbb{C}$ be non-zero and assume $a \neq \pm b$. A square root c of ab is called the *right choice* if

$$(7.27) \quad \left| \frac{a+b}{2} - c \right| \leq \left| \frac{a+b}{2} + c \right|.$$

It turns out that in order to have a limit for (1.1) one has to make the right choice for all but finitely many indices $n \geq 1$. This is described in detail by Cox [27].

7.9. Elliptic Landen with p -adic initial conditions. Let p be a prime and a, b be non-zero p -adic numbers. In order to guarantee that the p -adic series

$$(7.28) \quad c = a \sum_{i=0}^{\infty} \binom{\frac{1}{2}}{i} \left(\frac{b}{a} - 1\right)^i$$

converges, and thus defines a p -adic square root of ab , one must assume

$$(7.29) \quad b/a \equiv 1 \pmod{p^\alpha},$$

where $\alpha = 3$ for $p = 2$ and 1 otherwise. The corresponding sequence defined by (1.1) converges for $p \neq 2$ to a common limit: the p -adic AGM. In the case $p = 2$ one must assume that the initial conditions satisfy $b/a \equiv 1 \pmod{16}$. In the case $b/a \equiv 1 \pmod{8}$ but not 1 modulo 16, the corresponding sequence (a_n, b_n) does not converge, but the sequence of so-called *absolute invariants*

$$(7.30) \quad j_n = \frac{2^8(a_n^4 - a_n^2 b_n^2 + b_n^4)^3}{a_n^4 b_n^4 (a_n^2 - b_n^2)^2}$$

converges to a 2-adic integer. Information about these issues can be found in [32]. D. Kohel [36] has proposed a generalization of the AGM for elliptic curves over a field of characteristic $p \in \{2, 3, 5, 7, 13\}$. Mestre [45] has developed an AGM theory for ordinary hyperelliptic curves over a field of characteristic 2. This has been extended to non-hyperelliptic curves of genus 3 curves by C. Ritzenthaler [38]. An algorithm for counting points for ordinary elliptic curves over finite fields of characteristic $p > 2$ based on the AGM is presented in R. Carls [23].

7.10. Higher genus AGM. An algorithm analog to the AGM for abelian integrals of genus 2 was discussed by Richelot [47], [48] and Humboldt [34]. Some details are discussed by J. Bost and J. F. Mestre in [21]. The case of abelian integrals of genus 3 is due to D. Lehavi [38] and D. Lehavi and C. Ritzenthaler [39].

Gauss was correct: his numerical calculation (1.4) has grown in many unexpected directions.

Acknowledgements. The second author acknowledges the partial support of NSF-DMS0409968. The authors wish to thank Jon Borwein for many comments that lead to an improvement of the manuscript.

REFERENCES

- [1] M. Abramowitz and I. Stegun. *Handbook of Mathematical Functions with Formulas, Graphs and Mathematical Tables*. Dover, New York, 1972.
- [2] T. Amdeberhan, D. Manna, and V. Moll. The 2-adic valuation of a sequence arising from a rational integral. *Preprint*, 2007.
- [3] T. Amdeberhan and V. Moll. A formula for a quartic integral: a survey of old proofs and some new ones. *The Ramanujan Journal*, 2007.
- [4] B. Berndt and D. Bowman. Ramanujan's short unpublished manuscript on integrals and series related to Euler's constant. *Canadian Math. Soc. Conf. Proc.*, 27:19–27, 2000.
- [5] C. W. Borchardt. Über das arithmetisch-geometrische mittel aus vier elementen. *Berl. Monatsber.*, 53:611–621, 1876.
- [6] G. Boros and V. Moll. A criterion for unimodality. *Elec. Jour. Comb.*, 6:1–6, 1999.
- [7] G. Boros and V. Moll. An integral hidden in Gradshteyn and Ryzhik. *Jour. Comp. Applied Math.*, 106:361–368, 1999.
- [8] G. Boros and V. Moll. A rational Landen transformation. The case of degree 6. In Knopp G. Mendoza E.T. Quinto E. L. Grinberg S. Berhanu M, editor, *Contemporary Mathematics. Analysis, Geometry, Number Theory: The Mathematics of Leon Ehrenpreis*, volume 251, pages 83–89. American Mathematical Society, 2000.
- [9] G. Boros and V. Moll. The double square root, Jacobi polynomials and Ramanujan's master theorem. *Jour. Comp. Applied Math.*, 130:337–344, 2001.
- [10] G. Boros and V. Moll. Landen transformation and the integration of rational functions. *Math. Comp.*, 71:649–668, 2001.
- [11] G. Boros, V. Moll, and J. Shallit. The 2-adic valuation of the coefficients of a polynomial. *Scientia*, 7:37–50, 2001.
- [12] J. M. Borwein and D. H. Bailey. *Mathematics by Experiment: Plausible reasoning in the 21-st century*. A. K. Peters, 1st edition, 2003.
- [13] J. M. Borwein, D. H. Bailey, and R. Girgensohn. *Experimentation in Mathematics: Computational Paths to Discovery*. A. K. Peters, 1st edition, 2004.
- [14] J. M. Borwein and P. B. Borwein. The arithmetic-geometric mean and fast computation of elementary functions. *SIAM Review*, 26:351–366, 1984.
- [15] J. M. Borwein and P. B. Borwein. *Pi and the AGM- A study in analytic number theory and computational complexity*. Wiley, New York, 1st edition, 1987.
- [16] J. M. Borwein and P. B. Borwein. On the mean iteration $(a, b) \leftarrow ((a + 3b)/4, (\sqrt{ab} + b)/2)$. *Math. Comp.*, 53:311–326, 1989.
- [17] J. M. Borwein and P. B. Borwein. A remarkable cubic mean iteration. In St. Ruscheweyh, E.B. Saff, L. C. Salinas, and R. S. Varga, editors, *Computational Methods and Function Theory*, pages 27–31. Lectures Notes in Mathematics 1435, Springer-Verlag, 1990.
- [18] J. M. Borwein and P. B. Borwein. A cubic counterpart of Jacobi's identity and the AGM. *Trans. Amer. Math. Soc.*, 323:691–701, 1991.
- [19] J. M. Borwein, R. Crandall, and G. Fee. On the Ramanujan AGM fraction, I: the real parameter case. *Experimental Mathematics*, 13:275–285, 2004.
- [20] J. M. Borwein, R. Crandall, and G. Fee. On the Ramanujan AGM fraction, II: the complex-parameter case. *Experimental Mathematics*, 13:287–295, 2004.

- [21] J. B. Bost and J. F. Mestre. Moyenne arithmetico-geometrique et periodes des courbes de genre 1 et 2. *Gazette des Mathematiens, Soc. de Mathematique de France*, 38:36–64, 1988.
- [22] T. J. Bromwich. *An Introduction to the Theory of Infinite Series*. MacMillan, London, 2nd edition, 1926.
- [23] R. Carls. *A generalized arithmetic geometric mean*. PhD thesis, Gronigen, 2004.
- [24] B. Caviness and J. R. Johnson. *Quantifier elimination and cylindrical algebraic decomposition*. Texts and Monographs in Symbolic Computation. Springer-Verlag, 1st edition, 1998.
- [25] M. Chamberland and V. Moll. Dynamics of the degree six Landen transformation. *Discrete and Dynamical Systems*, 15:905–919, 2006.
- [26] G. E. Collins. Quantifier elimination for the elementary theory of real closed fields by cylindrical algebraic decomposition. *Lecture Notes in Computer Science*, 33:134–183, 1975.
- [27] D. Cox. The arithmetic-geometric mean of Gauss. *L'Enseignement Mathematique*, 30:275–330, 1984.
- [28] P. Deift. Continuous versions of some discrete maps or what goes on when the lights go out. *Jour. d'Analyse Math.*, 58:121–133, 1992.
- [29] K. F. Gauss. Arithmetisch Geometrisches Mittel. *Werke*, 3:361–432, 1799.
- [30] D. Grayson. The arithogeometric mean. *Arch. Math*, 52:507–512, 1989.
- [31] Kok Seng Chua Heng Huat Chan and P. Sole. Quadratic iterations to π associated with elliptic functions to the cubic and septic base. *Trans. Amer. Math. Soc.*, 355:1505–1520, 2002.
- [32] G. Henniart and J. F. Mestre. Moyenne arithmetico-geometrique p -adique. *C. R. Acad. Sci. Paris*, 308:391–395, 1989.
- [33] J. Hubbard and V. Moll. A geometric view of rational Landen transformation. *Bull. London Math. Soc.*, 35:293–301, 2003.
- [34] G. Humbert. Sur la transformation ordinaire des fonctions abeliennes. *Journal des Mathematiques*, 7, 1901.
- [35] M. Kauers and P. Paule. A computer proof of Moll's log-concavity conjecture. *Preprint*, 2007.
- [36] D. Kohel. The AGM- $X_0(N)$ Heegner point lifting algorithm and elliptic curve point counting. In *Proceedings of ASI-ACRYPT'03*, volume 2894 of *Lecture Notes in Computer Science*, pages 124–136. Springer-Verlag, 2003.
- [37] J. Landen. An investigation of a general theorem for finding the length of any arc of any conic hyperbola, by means of two elliptic arcs, with some other new and useful theorems deduced therefrom. *Philos. Trans. Royal Soc. London*, 65:283–289, 1775.
- [38] D. Lehavi. An explicit formula for the genus 3 AGM: arXiv:math.AG/011273.
- [39] D. Lehavi and C. Ritzenhaler. Formulas for the arithmetic geometric mean of curves of genus 3: arXiv:math.AG/0403182.
- [40] J. Little. On the zeroes of two families of polynomials arising from certain rational integrals. *Rocky Mountain Journal*, 35:1205–1216, 2005.
- [41] D. Manna and V. Moll. Rational Landen transformations on $\mathbb{R}$. *Math. Comp.*, 2007.
- [42] D. Manna and V. Moll. A simple example of a new class of Landen transformations. *Amer. Math. Monthly*, 114:232–241, 2007.
- [43] H. McKean and V. Moll. *Elliptic Curves: Function Theory, Geometry, Arithmetic*. Cambridge University Press, New York, 1997.
- [44] J. F. Mestre. Moyenne de Borchardt et integrales elliptiques. *C. R. Acad. Sci. Paris*, 313:273–276, 1991.
- [45] J. F. Mestre. Lettre adresse a Gaudry en Harley. 2000. <http://www.math.jussieu.fr/~mestre>.
- [46] D. J. Newman. A simplified version of the fast algorithm of Brent and Salamin. *Math. Comp.*, 44:207–210, 1985.
- [47] F. Richelot. Essai sur une methode generale pour determiner la valeur des integrales ultra-elliptiques, fondee sur des transformations remarquables de ces transcendentes. *C. R. Acad. Sci. Paris*, 2:622–627, 1836.
- [48] F. Richelot. De transformatione integralium Abelianorum primi ordinis commentation. *Jour. fur die reine und angew. Math.*, 16:221–341, 1837.
- [49] P. Sole. D_4 , E_6 , E_8 and the AGM. *Springer Lecture Notes in Computer Science*, 948:448–455, 1995.
- [50] P. Sole and P. Loyer. U_n lattices, construction B, and the AGM iterations. *Europ. J. Combinatorics*, 19:227–236, 1998.

- [51] G. Springer. *Introduction to Riemann Surfaces*. American Mathematical Society, 2nd edition, 2002.
- [52] G. N. Watson. The Marquis and the Land-agent. *Math. Gazette*, 17:5–17, 1933.
- [53] H. S. Wilf. *generatingfunctionology*. Academic Press, 1st edition, 1990.

DEPARTMENT OF MATHEMATICS AND STATISTICS, DALHOUSIE UNIVERSITY, HALIFAX, NOVA SCOTIA, CANADA, B3H 3J5

E-mail address: `dantemanna@gmail.com`

DEPARTMENT OF MATHEMATICS, TULANE UNIVERSITY, NEW ORLEANS, LA 70118

E-mail address: `vhm@math.tulane.edu`